

ERRATUM

Open Access



Erratum to: Glymphatic distribution of CSF-derived apoE into brain is isoform specific and suppressed during sleep deprivation

Thiyagaragan M. Acharyar^{1†}, Baoman Li^{1,4†}, Weiguo Peng¹, Philip B. Verghese², Yang Shi², Evan McConnell¹, Abdellatif Benraiss³, Tristan Kasper¹, Wei Song¹, Takahiro Takano¹, David M. Holtzman², Maiken Nedergaard¹ and Rashid Deane^{1*}

Erratum

The original article [1] contains an error whereby the co-author, Takahiro Takano's name was misspelt.

This error has now been corrected in the original article.

Author details

¹Center for Translational Neuromedicine, Division of Glial Disease and Therapeutics, Department of Neurosurgery, University of Rochester Medical Center, University of Rochester, Rochester, NY 14642, USA. ²Department of Neurology, Hope Center for Neurological Disorders, and the Charles F. and Joanne Knight Alzheimer's Disease Research Center, Washington University School of Medicine, St Louis, MO 63110, USA. ³Center for Translational Neuromedicine, Division of Cell and Gene Therapy, University of Rochester Medical Center, Rochester, NY 14642, USA. ⁴Laboratory of Brain Metabolic Diseases, Institute of Metabolic Disease Research and Drug Development, China Medical University, Shenyang, China.

Received: 23 December 2016 Accepted: 23 December 2016

Published online: 12 January 2017

References

1. Acharyar TM, et al. Glymphatic distribution of CSF-derived apoE into brain is isoform specific and suppressed during sleep deprivation. *Mol Neurodegener.* 2016;11:74. doi:10.1186/s13024-016-0138-8.

* Correspondence: Rashid_Deane@urmc.rochester.edu

[†]Equal contributors

¹Center for Translational Neuromedicine, Division of Glial Disease and Therapeutics, Department of Neurosurgery, University of Rochester Medical Center, University of Rochester, Rochester, NY 14642, USA

